



Energy transport: Species separation

- Separation of ion and electron energy fluxes requires determination of power exchange term.
 - At high collisionality ($n_e \sim 10^{20}$), this requires $O(10\text{eV})$ accuracy of $(T_e - T_i)$ profile, which has not yet been achieved.
 - Best analysis so far for highest T_i gives range from: **A)** large Q_e with $Q_i \sim Q_i^{\text{NC}}$ to **B)** $Q_i \sim Q_e \gg Q^{\text{NC}}$.
 - $Q_e \gg Q_i \sim Q_{\text{NC}}$ would be consistent with post-pellets experiments.
 - However, neoclassical electron energy fluxes *not* supported by measurements.
- > *Next campaign: Improvements in T_i profiles + heat wave measurements.*






Energy transport: Species separation

- Separation of ion and electron energy fluxes requires determination of power exchange term.
 - At high collisionality ($n_e \sim 10^{20}$), this requires $O(10\text{eV})$ accuracy of $(T_e - T_i)$ profile, which has not yet been achieved.
 - Best analysis so far for highest T_i gives range from: **A)** large Q_e with $Q_i \sim Q_i^{\text{NC}}$ to **B)** $Q_i \sim Q_e \gg Q^{\text{NC}}$.
 - $Q_e \gg Q_i \sim Q_{\text{NC}}$ would be consistent with post-pellets experiments.
 - However, neoclassical electron energy fluxes *not* supported by measurements.
- > *Next campaign: Improvements in T_i profiles + heat wave measurements.*






Energy transport: Species separation

- Separation of ion and electron energy fluxes requires determination of power exchange term.
 - At high collisionality ($n_e \sim 10^{20}$), this requires $O(10\text{eV})$ accuracy of $(T_e - T_i)$ profile, which has not yet been achieved.
 - Best analysis so far for highest T_i gives range from: **A)** large Q_e with $Q_i \sim Q_i^{\text{NC}}$ to **B)** $Q_i \sim Q_e \gg Q^{\text{NC}}$.
 - $Q_e \gg Q_i \sim Q_{\text{NC}}$ would be consistent with post-pellets experiments.
 - However, neoclassical electron energy fluxes *not* supported by measurements.
- > Next campaign: Improvements in T_i profiles + heat wave measurements.






Energy transport: Species separation

- Separation of ion and electron energy fluxes requires determination of power exchange term.
 - At high collisionality ($n_e \sim 10^{20}$), this requires $O(10\text{eV})$ accuracy of $(T_e - T_i)$ profile, which has not yet been achieved.
 - Best analysis so far for highest T_i gives range from: **A)** large Q_e with $Q_i \sim Q_i^{\text{NC}}$ to **B)** $Q_i \sim Q_e \gg Q^{\text{NC}}$.
 - $Q_e \gg Q_i \sim Q_{\text{NC}}$ would be consistent with post-pellets experiments.
 - However, neoclassical electron energy fluxes *not* supported by measurements.
- > Next campaign: Improvements in T_i profiles + heat wave measurements.






Energy transport: Species separation

- Separation of ion and electron energy fluxes requires determination of power exchange term.
 - At high collisionality ($n_e \sim 10^{20}$), this requires $O(10\text{eV})$ accuracy of $(T_e - T_i)$ profile, which has not yet been achieved.
 - Best analysis so far for highest T_i gives range from: **A)** large Q_e with $Q_i \sim Q_i^{\text{NC}}$ to **B)** $Q_i \sim Q_e \gg Q^{\text{NC}}$.
 - $Q_e \gg Q_i \sim Q_{\text{NC}}$ would be consistent with post-pellets experiments.
 - However, neoclassical electron energy fluxes *not* supported by measurements.
- > Next campaign: Improvements in T_i profiles + heat wave measurements.



Energy transport: Species separation

- Separation of ion and electron energy fluxes requires determination of power exchange term.
 - At high collisionality ($n_e \sim 10^{20}$), this requires $O(10\text{eV})$ accuracy of $(T_e - T_i)$ profile, which has not yet been achieved.
 - Best analysis so far for highest T_i gives range from: **A)** large Q_e with $Q_i \sim Q_i^{\text{NC}}$ to **B)** $Q_i \sim Q_e \gg Q^{\text{NC}}$.
 - $Q_e \gg Q_i \sim Q_{\text{NC}}$ would be consistent with post-pellets experiments.
 - However, neoclassical electron energy fluxes *not* supported by measurements.
- > Next campaign: Improvements in T_i profiles + heat wave measurements.



Energy transport: Species separation

- Separation of ion and electron energy fluxes requires determination of power exchange term.
 - At high collisionality ($n_e \sim 10^{20}$), this requires $O(10\text{eV})$ accuracy of $(T_e - T_i)$ profile, which has not yet been achieved.
 - Best analysis so far for highest T_i gives range from: **A)** large Q_e with $Q_i \sim Q_i^{\text{NC}}$ to **B)** $Q_i \sim Q_e \gg Q^{\text{NC}}$.
 - $Q_e \gg Q_i \sim Q_{\text{NC}}$ would be consistent with post-pellets experiments.
 - However, neoclassical electron energy fluxes *not* supported by measurements.
- > Next campaign: Improvements in T_i profiles + heat wave measurements.

